SEUREI
Approved For Release 2001/07107 [CNA] APPROVED FOR RELEASE 2001

6-2429 6 August 1967

## Eumary

North Vietnesses army troops infiltrating into South Vietness through Loss during 1966 apparently sustained permanent losses of about 20 percent.\* At least 53,000 and possibly 81,000 EVA troops are estimated to have successfully infiltrated into South Vietness during 1966. Thus, from 10,000 to 15,000 men probably were permanently lost infiltrating during 1966. Three fourths of these losses resulted from death and personent incapacitation due to illness -- overwhelmingly maleria. About ten percent of those lost were killed by air attacks and about five percent are considered permanent deserters. The loss rate for 1966 appears to have increased perhaps several times over the 1965 rate. To data are available for 1967. Increased incidence of serious eickness was responsible for most of the higher 1966 rate. In addition, although there were no losses found in the reports surveyed which were directly attributable to air strikes during 1965, losses from this cause in 1966 were about ten percent of the total. Perhaps as many as one half of all infiltrating troops suffer to some degree from maleria en route to fouth Vietnem. As a result

NO FOREIGN DISSEM
SECRET

<sup>\*</sup> These estimates are based on a limited sample of data on infiltration and should be considered preliminary. See methodology below. Infiltration through the DMZ is excluded.

of this and other known health problems perhaps as many men are not fit for combat when they arrive as are lost en route. Nost of these men probably recover sufficiently for combat.

Infiltration through Laos

South Vietness runs through the eastern portion of the Lactian penhandle. Troops move from their point of origin in the North to the general vicinity of Hill 1901 just north of the extreme western part of the INZ. In the beginning all troops were trucked to the Hill 1901 area, but during 1966 were increasingly forced to move on foot as a result of air interdiction in the southern areas of North Vietness. A series of trails leads south from Hill 1901 through the western corner of the INZ into Laos, and from there along the Laos/SVN border and enters South Vietness at Quang Tri. Thus Thien, Quang Nam and other provinces south. Depending on the point of entry into SVN, the trip through Leos takes between one and four months.

The 20 percent estimate calculated for permanent losses from all causes in 1966 is supported by the interrogation of one senior ECO from a regimental size unit who stated that infiltration units usually are from 10 to 30 percent above authorized strength to compensate for attrition.

Although air interdiction apparently does not cause many direct casualties, it has had other significant effects. Units have been forced to change their mode of movement in EVS from truck to travel on foot increasing the time needed to infiltrate. (All infiltration movement in

-0-

NO FOREIGN DISSEM

Leos has been by foot.) The additional time spent travelling by foot in MVN results in a higher rate of sickness in MVN and also in Isos.

During 1966, at least 53,000 and possibly 81,000 MVA troops were estimated to have infiltrated into South Vietnem. If 75 percent of them came through Leos, then based on the 20 percent permanent loss rate, 10,000 and possibly 15,000 men were lost while infiltrating. (.75 x 53,000 = 39,750. .80 x = 59,750 = 49,687. 49,687 infiltrated, 9,937 were lost, and 39,750 arrived.)

## Methodology

The table below summarizes the data used for the estimates in this report. Evidence of about 7,000 infiltrators in eleven units varying from company to regiment in size make up the data base. Four units with 1,777 men infiltrated during the last half of 1965 and seven units with 5,390 men infiltrated throughout 1966. Information was taken from all interrogation reports is mediately svailable. Four interrogation methods caused gaps in information and as a result the data base for each estimate is less than the total. To produce estimates of permanent losses it is assumed on the basis of limited evidence that one balf of all deserters eventually are returned to duty and that two-thirds of the men who drop out of their units during infiltration are eventually returned to duty.

-2-

NO FOREIGN DISSEM SECRE

		Arriving in SVN during last Arriving in SVN half of 1965 during 1966
(1)	Deserters	14 of 1277 - 1 percent 60 of 2890, - 2 percent
(2)	Killed by Air Attack	0 of 200 = 0 percent 94 of 4270 = 2 percent
(3)	Died from Sickness	23 of 1407 = 2 percent 79 of 1270 = 6 percent
(4)	Propped out of Unit*	42 of 1277 - 3 percent 117 of 330 - 30 percent
	For 1965: 1/2 of (1)	plus 1/3 of (4) plus (2) and (3) = 3.5 percent.
For	1966: 1/2 of (1) plus	1/3 of (4) plus (2) and (3) = 19 percent.

<sup>\*</sup> Although some portion of this sharp increase may result from the inadequate data base used, there are reasons to believe that there has been a steady increase in sickness. Troops that march in NVN to Lace begin their travel in Lace in a weakened condition and medical care in Lace has deteriorated.

Distributions

Orig. & 1 - Addressee

1 - D/OER

1 - ADD/OER

1 - St/PR

T = 20/C

T - DAVA

T - D/T

1 - 1/8

rew/9326-R (8 Aug 67)

25X1A

NO FOREIGN DISSEM SECRET